

1 **Investigating the status of pyrethroid resistance in UK populations of the**
2 **cabbage stem flea beetle (*Psylliodes chrysocephala*)**

3

4 Caitlin Willis^{a,d}, Stephen P. Foster^a, Christoph T. Zimmer^b, Jan Elias^b, Ralf Nauen^c, Xianmin
5 Chang^d, Linda M. Field, Martin S. Williamson^a & T.G. Emyr Davies^{a*}

6 ^aBiointeractions and Crop Protection Department, Rothamsted Research, West Common,
7 Harpenden, AL5 2JQ, UK.

8 ^bSyngenta Crop Protection, Werk Stein, Schaffhauserstrasse, Stein CH4332, Switzerland

9 ^cBayer AG, Crop Science Division, R&D, Monheim, Germany.

10 ^dRoyal Agricultural University, Stroud Rd, Cirencester, Gloucestershire, UK.

11

12 ***Corresponding author:** E-mail address: emyr.davies@rothamsted.ac.uk (T.G. Emyr Davies)

13 **Declaration of interest:** none

14

HIGHLIGHTS

15 • UK populations of cabbage stem flea beetle exhibit high levels of resistance to pyrethroid
16 insecticides

17 • Pyrethroid resistance in UK populations is largely the result of increased metabolism

18 • Resistance levels are highest in the South East of the UK

19 • Resistance is advancing to the North and West of England.

20

21 **ABSTRACT**

22 The cabbage stem flea beetle, *Psylliodes chrysocephala* L. is a major pest of winter oilseed
23 rape in several European countries. Traditionally, pyrethroid insecticides have been widely
24 used for control of *P. chrysocephala*, but in recent years control failures have occurred. In line
25 with previous surveys, UK populations of *P. chrysocephala* were found to exhibit high levels
26 of resistance to the pyrethroid lambda-cyhalothrin. This resistance was suppressed by pre-
27 treatment with the cytochrome P450 inhibitor PBO, suggesting that the resistance remains

28 largely metabolic. The L1014F (kdr) mutation in the voltage-gated sodium channel, which
29 confers relatively low levels (10-20 fold) of resistance to pyrethroids, was also found to be
30 widespread across the UK regions sampled, whereas the L925I (super-kdr) mutation was also
31 present but much less common. The current survey also suggests that higher levels of
32 pyrethroid resistance are starting to spread to the North and West of England, and that
33 resistance levels continue to remain high in the South East.

34

35 **Keywords:** cabbage stem flea beetle; oilseed rape; pyrethroid resistance

36 1. Introduction

37 The cabbage stem flea beetle, *Psylliodes chrysocephala* (Coleoptera: Chrysomelidae) is an
38 established and key insect pest of winter oilseed rape, particularly in the UK (Graham and
39 Alford, 1981) and Germany (Zimmer *et al.*, 2014), and is a significant pest of other *Brassica*
40 species in several European countries (Bromand, 1990; Bartlet and Williams, 1991; Bartlet,
41 Mithen and Clark, 1996). *P. chrysocephala* inflicts damage at both the larval and adult stage,
42 with the tunnelling of the larvae into the leaf petioles and main stems causing the most damage
43 through weakening of the upper section of the roots and lower parts of the stems (Williams,
44 2004). When infestation is high, the plant tips distort, the stems wilt and the infested plants
45 become more susceptible to fungal infections such as *Phoma lingam* (Alford, 2003), the
46 bacterial disease *Erwinia* sp. and frost damage (Højland *et al.*, 2015; Højland and Kristensen,
47 2018). Adult *P. chrysocephala* cause damage by feeding on stems, cotyledons and the first true
48 leaves during crop emergence resulting in ‘shot-holing’ symptoms, leading to poor plant vigour
49 or potential seedling death before emergence when fields are heavily infested (Zimmer *et al.*,
50 2014). Prior to 2014, *P. chrysocephala* affected approximately 67% of the area of oil seed rape
51 grown in the UK causing an annual 1% yield loss (Clarke *et al.*, 2009). However, in 2014,
52 serious crop losses due to adult beetles (2.7% of the national crop) were recorded, the most
53 serious losses (5–14%) being in eastern and southern England (Wynn, Ellis and Alves, 2014).
54 In the autumn of 2015, a more extensive survey found that over 65% of crops had some
55 damage, and that the damage was more widely distributed across the country than in 2014,
56 although nationally only 1% of crops was lost (Alves, Wynn and Stopps, 2015). Subsequent
57 surveys have confirmed that the average numbers of larvae per plant have risen substantially
58 in all regions since 2014 (as summarised in Dewar, 2017).

60 Prior to December 2013, control of *P. chrysocephala* relied on the protection of oilseed rape
61 seedlings by systemic neonicotinoid seed treatments containing either imidacloprid,
62 thiamethoxam or clothianidin, followed by the application of foliar pyrethroid sprays later in
63 the season if needed (Højland *et al.*, 2015; Højland and Kristensen, 2018). However, in
64 December 2013, the European regulatory authorities (EU commission, 2013) banned the use
65 of neonicotinoid seed treatments on all outdoor flowering crops, thus preventing their use in
66 oil seed rape, leading to the increase in *P. chrysocephala* and the increased use of pyrethroid
67 sprays. Today, pyrethroids (e.g. lambda-cyhalothrin) are the only class of insecticide that
68 remain for chemical control of *P. chrysocephala* in the UK and other parts of mainland Europe.

69 The continuous use of pyrethroids to control *P. chrysocephala*, coupled with the lack of
70 alternative insecticides with different modes of action, has led to a high selection pressure,
71 driving the development and spread of resistance. Resistance to lambda-cyhalothrin was first
72 reported in 2008, in north-western Mecklenburg, Western Pomerania, a major oilseed rape
73 growing area in Northern Germany (Heimbach and Müller, 2013). Zimmer *et al.*, (2014)
74 reported the presence of the L1014F *kdr* mutation in the voltage-gated sodium channel, with
75 high frequencies of the allele (90-100%) being found in populations collected from across
76 Northern Germany, with the beetles exhibiting a low level resistance against a range of
77 pyrethroids including lambda-cyhalothrin and tau-fluvalinate. More recently, studies by
78 Højland *et al.*, (2015) and Højland and Kristensen (2018) have shown that pyrethroid resistance
79 resulting from the *kdr* mutation has continued to spread northwards, and is now present in
80 populations from both Denmark and the UK, whilst in Germany it has spread further south.

81 Despite the presence of *kdr* in UK populations, Højland *et al.*, (2015) found that the high
82 pyrethroid resistance levels, with control failures being observed at the full field rate, did not
83 completely correlate with the *kdr* genotype suggesting that another mechanism of resistance,
84 such as metabolic resistance, is also present. Given the lack of alternative insecticides with

85 different modes of action, the presence and spread of pyrethroid resistance is concerning for
86 the chemical control of *P. chrysocephala*.

87 The present study has determined the current status, extent and geographical spread of
88 pyrethroid resistance in UK populations of *P. chrysocephala*. Bioassays, based on glass vial
89 exposure of adult beetles to lambda-cyhalothrin, were carried out on samples collected in 2018
90 and 2019 to examine how resistance had changed over this time across the UK. The presence
91 of the *kdr* and super-*kdr* target-site mutations in UK populations was also monitored, and the
92 potential contribution of a metabolic resistance component in the beetles assessed by pre-
93 treatment with the synergist PBO, which is a cytochrome P450 inhibitor.

94 **2. Methods**

95 2.1 Collection of field samples of *Psylliodes chrysocephala*

96 In July/August 2018 and 2019, live *P. chrysocephala* adults were collected from oil seed rape
97 pods freshly harvested from the fields at Rothamsted Research, Harpenden, Hertfordshire,
98 using a hand-held battery-powered pooter. Insects were maintained at 15±1°C, with 65%
99 relative humidity in a light:dark photoperiod of 12:12h. Adults were kept in a mesh cage and
100 fed continuously on a diet of Chinese cabbage (*Brassica rapa* spp). Further samples were
101 received by post from oilseed rape fields across the UK and were kept in sealed plastic bags or
102 plastic containers containing Chinese cabbage or oilseed rape plant material and moist tissue
103 paper, maintained in the same environmental conditions as the Rothamsted samples.

104 2.2 Bioassays to test the effect of pyrethroids on *Psylliodes chrysocephala*

105 *P. chrysocephala* samples were tested for resistance to the pyrethroid lambda-cyhalothrin
106 (Syngenta, UK) using a glass vial bioassay based on IRAC (Insecticide Resistance Action
107 Committee) Method 031 (www.irac-online.org/methods/weevils-and-flee-beetles/2014). Glass
108 vials (14ml: 7cm tall/ 2cm diameter) (S Murray and Co, Surrey, UK) were prepared by coating

the inner surface with different concentrations of the insecticide. Initial stock solutions were prepared by diluting the technical grade insecticide in technical grade acetone. Three doses, equivalent to 4%, 20% and 100% of the recommended field application rate of lambda-cyhalothrin (7.5 g ai/ha) were used (Table 1). The controls were glass vials treated with acetone only. To coat vials, 500µl of solution was pipetted into the vials which were then placed horizontally without lids on a roller in a fume hood. Vials were rotated at room temperature for at least 2 hours until all the acetone had evaporated. Vials were then left vertically at 4°C overnight before attaching the screw tops the following day.

Table 1. Concentrations of lambda-cyhalothrin tested on *P. chrysocephala* via glass vial bioassay.

Concentration		% Field rate
(g/ha)	ppm	
0.3	0.204	4
1.5	1.02	20
7.5	5.1	100

The adult beetles (see 2.1) were used within a few days of collection and only those capable of walking or jumping when released onto a tray inside a three-sided Perspex cage were collected, using a hand-held battery-powered pooter. A minimum of ten beetles were transferred from the inverted pooter through a small funnel into each vial. The vials were then resealed and left at 18±1°C under a 16:8h light:dark photoperiod. After 24 hours, the beetles were transferred to untreated glass vials without lids under upturned 200ml plastic disposable cups (VWR International Ltd, Dublin, Ireland), to allow for a potential recovery which can occur in insects with metabolic resistance. After a further 24 hours, the beetles were released onto a tray and individuals scored using a fine paint brush according to three categories: ‘mobile’ (capable of jumping or walking in a coordinated way), ‘affected’ (incapable of jumping or coordinated movement) or ‘dead’ (no movement). Scoring of the beetles from each vial was done for 10 minutes to avoid adults that were simulating death, a behaviour shown by this species that has

probably evolved through predation pressure. Results were expressed as percentage mortalities. Following scoring, beetles in each category were transferred to Eppendorf tubes and snap frozen using liquid nitrogen before being stored in a freezer at -80°C.

2.3 TaqMan PCR assay to detect the presence of kdr/skdr in *Psylliodes chrysocephala*

TaqMan genotyping assays (Livak, 1999) were used to determine the presence of the mutations responsible for the kdr (L1014F) and super-kdr (L925I) sodium channel substitutions in individual adult beetles. Primer Express v.2.0 (Life Technologies) was used to design the primer and probe sequences for the assays (Table 2). In both assays, VIC reporter dye-labelled probes were used to detect the wild-type susceptible allele and 6-FAM reporter dye-labelled probes to detect the resistant allele. Each probe contained a 3' non-fluorescent quencher dye.

Table 2. Primer and probe sequences used for TaqMan assays to detect the L1014F (kdr) and L925I (skdr) mutations in *Psylliodes chrysocephala*.

Primer/Probe		Sequence	144
Primers	kdr-F	GGACTGTATGCTAGTCGGTGATGT	145
	kdr-R	GCAAAGCCAAGAAGAGATTTCAGTA	
	skdr-F	GCCAAGTCATGGCCAACTT	146
	skdr-R	TATAATGCACAGCACAAAGGTCA	
Probes	kdr-VIC	TTACCACAAGATTACC	147
	kdr-FAM	TTACCACAAAATTACC	
	skdr-VIC	TGGGTGCTTTAGGTAA	148
	skdr-FAM	TGGGTGCTATAGGTAA	
			149

PCR reactions (15µl) contained 1.5µl (50ng) genomic DNA, 7.5µl SensiFast probe mix (Bioline Reagents Ltd, UK), 0.375µl of kdr or skdr primer/probe mix (800nM of each primer and 200nM of each probe) and sterile water. Reactions were run on an Applied Biosystems 7900HT real-time PCR system, with initial incubations at 50°C for 2 minutes and 95°C for 10 minutes, followed by 40 cycles of 95°C for 15 seconds and 60°C for 45 seconds. The increase

in VIC and 6-FAM reporter dye fluorescence was monitored in real time and an allelic discrimination analysis performed using the 7900HT Sequence Detection System software.

2.4 Use of a synergist to identify the presence of metabolic resistance in *Psylliodes chrysocephala*

Pre-treatment with the insecticide synergist Piperonyl butoxide (PBO), obtained from Sigma-Aldrich (Missouri, USA), was used to detect potential metabolic resistance mechanisms *in-vivo*. PBO was diluted in technical grade acetone to give an equivalent concentration of 4g aiha⁻¹. 500µl of solution was then used to coat glass vials (see 2.2). Ten beetles per replicate were transferred to the PBO-coated vials for 1 hour before being transferred to either untreated control vials or vials coated with lambda-cyhalothrin at the 100% field rate (7.5 g aiha⁻¹). The beetles were then bio-assayed in parallel to beetles from the same sample not pre-exposed to PBO.

3. Results and Discussion


3.1 Survey of pyrethroid resistance in *Psylliodes chrysocephala* across the UK

To determine the current extent and geographical spread of resistance to pyrethroid insecticides in UK *P. chrysocephala* populations, and how this compares to previous reports (Højland *et al.*, 2015), bioassays with lambda-cyhalothrin were conducted on adult beetle samples from Rothamsted Research's farm in Hertfordshire and oilseed rape fields located across England, Scotland and Wales. The bioassays allowed the samples to be categorised as being either completely susceptible, or to contain beetles that were 0-25%, 25-50%, 50-75%, 75-99% and 100% resistant, depending on the percentage of beetles per sample surviving treatment with 7.5 g ai ha⁻¹ lambda-cyhalothrin. The bioassay is an approved test method (method 031) for determining resistance in *P. chrysocephala* (IRAC, 2014) and was used by Zimmer *et al.*, (2014) to monitor the emergence and geographic spread of pyrethroid resistance in *P.*

chrysocephala in Germany, by Højland *et al.*, (2015) to determine the spread of pyrethroid resistance in Danish, British and German samples and most recently by Højland and Kristensen (2018) when investigating lambda-cyhalothrin resistance in Danish populations. Similar bioassays have also been used to monitor the spread of pyrethroid resistance in European populations of pollen beetle (*Meligethes aeneus*), another major pest of oilseed rape (Zimmer and Nauen (2011), Slater *et al.*, (2011), Nauen *et al.*, (2012)).

In 2018, a total of 42 *P. chrysocephala* samples, obtained from four different regions across England, but primarily from counties in the East (Fig. 1), were tested. Of these only five samples were found to contain no mobile beetles at 100% of the recommended field rate for lambda-cyhalothrin, which would be expected if the sample was susceptibility. However, for these five samples mortality was found to be <90% at 20% of the field rate, suggesting resistance is present as judged by the IRACs 'susceptibility rating scheme' (IRAC, 2014). The other 37 samples all showed some level of resistance with the highest resistance, at 89% being the sample from Bishop Cannings (Wiltshire).

In 2019 a total of 145 *P. chrysocephala* samples were obtained from across England, representing more of the country (Fig. 1), two samples were received from Wales and one from Scotland. Only the Scottish sample was found to be truly susceptible to lambda-cyhalothrin, displaying 100% mortality at 20% of the recommended field rate. Worryingly, populations containing 100% resistant beetles were recorded for the first time in the UK. Overall, the distribution maps for pyrethroid resistance in UK populations of *P. chrysocephala* (Fig. 1) suggest that higher levels of resistance are starting to spread to the North and West of England and that resistance levels continue to remain high in the South East.



213

2018

214

215 **Figure 1.** Pyrethroid resistance in *P. chrysocephala* in the UK for 2018 and 2019. The maps
 216 were created using QGIS (version 3.0.3) and use a 6-category colour scale to show the level of
 217 resistance. The map is divided into counties (light grey borders) and regions (dark grey
 218 borders).

219

220 Over the two years of monitoring, the percentage of pyrethroid-resistant beetles in each sample
 221 increased (Fig. 2). In 2018 the percentage in the 0-10% resistance category was approx. 16%
 222 whereas in 2019 this was down to 5%. In 2018 there were no samples in the 90-100% resistance
 223 category whereas in 2019 this increased to approx. 5%. The median resistance value was 28.5%
 224 in 2018 and 60% in 2019.

225

226





Figure 2. Histograms showing the shift in the relative % frequency of pyrethroid resistant *P. chrysocephala* in 2018 and 2019.

The median resistance levels recorded in 2019 were highest (approx. 60%) in the East Midlands, South East, South West and Yorkshire, and the Humber (Fig. 3). However, overall in each region the resistance levels were fairly evenly spread, with no statistically significant difference in resistance being found between regions.




Figure 3. Boxplot of % resistance to pyrethroid in *P. chrysocephala* in each region of the UK surveyed in 2019.

When these 2019 data are broken down further to county level (Fig. 4), it is clear that there are also no significant differences in resistance across counties and that there are no resistance ‘hotspots’. This suggests resistance is highly localised, almost on a farm-by-farm basis.




Figure 4. Boxplot of % resistance to pyrethroid in *P. chrysocephala* in each county of the UK surveyed in 2019.

3.2 Pyrethroid resistance mechanism(s) in *P. chrysocephala*

The TaqMan assays (see 2.3) were used to detect the presence of the L1014F (*kdr*) and L925I (*s-kdr*-like) substitutions in the 2018 and 2019 *P. chrysocephala* samples (Table 3). In 2018, 40 beetles from seven UK samples of *P. chrysocephala* were tested using only individuals that survived the 100% field rate of lambda-cyhalothrin, enabling the genotype associated with the resistant, mobile phenotype to be determined. The samples were from Great Saxham (Suffolk), Bishop Cannings (Wiltshire), Rothamsted (Hertfordshire), Linton (Cambridgeshire), Feltwell (Norfolk) and Horbling and Grantham (Lincolnshire). The L1014F mutation was present at all of the sites, with 47.5% of the beetles being homozygous for the resistant allele (RR), 37.5% heterozygous (SR) and the remaining 15% *kdr* SS, although this genotype was not present in the Suffolk or Wiltshire populations. In the Lincolnshire, Cambridge, Norfolk and Hertfordshire samples the *kdr* SS genotype represented 5%, 5%, 2.5% and 2.5% of the mobile beetles respectively. The detection of *kdr* SS genotypes in beetles that displayed the mobile phenotype after treatment with the label rate of lambda-cyhalothrin, suggests the presence of

another resistance mechanism in *P. chrysocephala*. In contrast to L1014F, the L925I mutation was much less common, with two samples (Norfolk and Rothamsted) being all SS and the overall percentage of beetles showing the homozygous L925I genotype (RR) being only 2.5%. This suggests that a potential fitness cost is associated with this mutation.

In 2019, *P. chrysocephala* individuals were screened for *kdr* from sites close to those sampled in 2018 (although a sample from Oxfordshire was also included) and again, only beetles that survived the 100% field rate of lambda-cyhalothrin were used. The L1014F mutation was present at all sites except the one from Scotland. The percentage of beetles homozygous for the *kdr* resistance allele (RR) increased in three of the samples, Suffolk, Norfolk and Hertfordshire but decreased overall from 47.5% to 36%. Given that the percentage of beetles resistant to lambda-cyhalothrin in each sample increased between 2018 and 2019, but there was an overall decrease in the homozygous and heterozygous L1014F mutation, this further suggests the presence of another resistance mechanism in *P. chrysocephala*. Whilst the L925I mutation was less common than the L1014F mutation, it was found to be present in the Wiltshire and Hertfordshire samples which contained only the wild-type metabolic genotype (SS) in 2018. In the Oxford sample 15% of the beetles tested for the s-*kdr* mutation were homozygous for the resistant allele (RR). Overall the percentage of beetles showing the homozygous genotype (RR) was 6.6%.

2018

Region	County	Populations	No°	<i>kdr status</i>			
				SS	SR	RR	%RR
East Midlands	Suffolk	1	4	0	3	1	25%
	Wiltshire	1	8	0	4	4	50%
	Lincolnshire	2	10	2	2	6	60%
East of England	Cambridgeshire	1	8	2	2	4	50%
	Norfolk	1	6	1	3	2	33%
	Hertfordshire	1	4	1	1	2	50%
	Total	7	40	6 (15%)	15 (37.5%)	19 (47.5%)	

Region	County	Populations	No°	<i>skdr status</i>			
				SS	SR	RR	%RR
East Midlands	Suffolk	1	4	3	1	0	0%
	Wiltshire	1	8	5	3	0	0%
	Lincolnshire	2	10	9	1	0	0%
East of England	Cambridgeshire	1	8	5	2	1	13%
	Norfolk	1	6	6	0	0	0%
	Hertfordshire	1	4	4	0	0	0%
	Total	7	40	32 (80%)	7 (17.5%)	1 (2.5%)	

296

297 **2019**

Region	County	Populations	No°	<i>kdr status</i>			
				SS	SR	RR	%RR
East Midlands	Suffolk	1	4	0	1	3	75%
	Wiltshire	1	5	3	1	1	20%
	Lincolnshire	2	16	2	8	6	38%
East of England	Cambridgeshire	1	8	1	6	1	13%
	Norfolk	1	8	2	4	5	63%
	Hertfordshire	1	5	1	1	3	60%
SE England	Oxfordshire	1	19	7	4	8	42%
Scotland	Aberdeenshire	1	10	10	0	0	0%
	Total	9	75	26 (34.7%)	25 (33.3%)	27 (36%)	

298

Region	County	Populations	No°	<i>skdr status</i>			
				SS	SR	RR	%RR
East Midlands	Suffolk	1	5	5	0	0	0%
	Wiltshire	1	3	1	1	1	33%
	Lincolnshire	2	16	11	5	0	0%
East of England	Cambridgeshire	1	8	5	3	0	0%
	Norfolk	1	8	7	1	0	0%
	Hertfordshire	1	6	4	1	1	17%
SE England	Oxfordshire	1	20	13	4	3	15%
Scotland	Aberdeenshire	1	10	10	0	0	0%
	Total	5	76	56 (73.7%)	15 (19.7%)	5 (6.6%)	

299

300 **Table 3.** Detection of *kdr/skdr* alleles in *P. chryscephala* using TaqMan assay

301

302

303

3.3 Bioassays of *P. chrysocephala* using lambda-cyhalothrin and the synergist PBO

The insecticide synergist piperonyl butoxide has been shown to inhibit both P450 monooxygenases and esterases, thereby acting as a tool for the identification of metabolic resistance in insect samples (Young, Gunning and Moores, 2006). To investigate the lack of correlation between lambda-cyhalothrin resistance and *kdr* frequency, and to determine whether P450 monooxygenases (and or esterases) may play a role in mediating pyrethroid resistance in UK *P. chrysocephala* populations, synergist bioassays with PBO pre-treatments were conducted on five *P. chrysocephala* samples.

When exposed to lambda-cyhalothrin at the recommended field rate, the percentage of beetles affected was 47% (North Yorkshire), 75% (Wiltshire), 8% (Wiltshire), 40% (Leicestershire) and 40% (Hertfordshire) (Fig. 5). However, all adults pre-treated with PBO, prior to exposure to lambda-cyhalothrin at the same field rate were killed. This strongly suggests that a metabolic-based mechanism for pyrethroid resistance is present in the *P. chrysocephala*.




Figure 5. Restoration of insecticide (pyrethroid) susceptibility in *P. chrysocephala* following pre-treatment with PBO. Samples tested were from North Yorkshire, Wiltshire (x2), Leicestershire and Hertfordshire.

4. Conclusions

Since the EU-imposed ban on neonicotinoid seed treatments, the pyrethroid insecticide lambda-cyhalothrin has been widely used for chemical control of *P. chrysocephala* in the UK. This has resulted in a high selection pressure and led to the development of resistance, particularly in the South East of England. In the current study, populations of *P. chrysocephala* from around the UK were found to exhibit high levels of resistance to lambda-cyhalothrin, but some of this resistance was suppressed by the cytochrome P450 inhibitor PBO. This suggests that, as well as target site resistance, there is P450 mediated- detoxification of lambda-cyhalothrin, although further research is required to identify the specific P450(s) involved and elucidate the exact mechanism of resistance. This resistance to pyrethroids has resulted in widely-reported control problems for this pest in the farming press (e.g. Clark 2014; Casswell, 2014; FarmingUK team, 2015; Hill, 2017; FarmingUK team, 2017; Case, 2018; Allison, 2019; Dyer, 2019; Gillbard, 2019) since the introduction of the neonicotinoid ban.

The development of pyrethroid resistance in *P. chrysocephala* is a matter of growing concern, with the increasing spread of resistance threatening the production of oilseed rape both in the UK and mainland Europe (Heimbach and Müller, 2013). Thus, continuing to monitor the extent and geographical spread of pyrethroid resistance in this pest is particularly important at a time when synthetic pesticides are becoming less effective, through the evolution and selection of resistance, and less favoured, through EU legislation. Clearly there needs to be informed decision making on how to best deploy pesticides effectively in the future.

Acknowledgements

This work was funded by the John Oldacre Foundation (registered charity number 284960) through a PhD scholarship awarded to CW and by a Syngenta Crop Protection CASE award. The Smart Crop Protection (SCP) strategic programme (BBS/OS/CP/000001) at Rothamsted

Research is funded through the Biotechnology and Biological Sciences Research Council's Industrial Strategy Challenge Fund.

References

Alford, D. (2003) *Biocontrol of Oilseed Rape Pests*. John Wiley & Sons.

Allison, R. (2019). Cabbage stem flea beetle losses threaten oilseed rape's place in the rotation. *Farmers Weekly*. [online] Available at: <https://www.fwi.co.uk/arable/crop-management/pests/more-than-half-of-osr-plants-infested-with-flea-beetle-larvae> [Accessed 29 Jan. 2020].

Alves L, Wynn S and Stopps J, Cabbage stem flea beetle live incidence and severity monitoring 2015. AHDB Project Report No. 551. [Online]. Available: <http://cereals.ahdb.org.uk/media/799228/pr551-final-project-report.pdf>

Barres, B., Micoud, A., Corio-Costet, M.F., Debieu, D., Fillinger-David, S., Walker, A.S., Délye, C., Grosman, J. and Siegwart, M. (2016) Trends and challenges in pesticide resistance detection. *Trends in Plant Science*, 21(10), pp.834-853.

Bartlet, E., Mithen, R. and Clark, S. J. (1996) Feeding of the cabbage stem flea beetle *Psylliodes chrysocephala* on high and low glucosinolate cultivars of oilseed rape, *Entomologia Experimentalis et Applicata*. Kluwer Academic Publishers.

Bartlet, E. and Williams, I. H. (1991) 'Factors restricting the feeding of the cabbage stem flea beetle (*Psylliodes chrysocephala*)', *Entomologia Experimentalis et Applicata*.

Wiley/Blackwell (10.1111), 60(3), pp. 233–238.

Bromand, B. (1990) Diversities in oilseed rape growing in the Western Palaearctic region. OILB/WPRS Bulletin 13(4): 7–31.

Case, P. (2018). Farmers report severe flea beetle attacks in emerging oilseed rape crops.

378 *Farmers Weekly*. [online] Available at: <https://www.fwi.co.uk/arable/crop->
379 [management/pests/farmers-report-severe-flea-beetle-attacks-in-emerging-oilseed-rape-crops](https://www.fwi.co.uk/arable/crop-management/pests/farmers-report-severe-flea-beetle-attacks-in-emerging-oilseed-rape-crops)
380 [Accessed 10 Feb. 2020].

381 Caswell, L. (2014). OSR growers face disaster from flea beetle attack. *Farmers Weekly*.
382 [online] Available at: [https://www.fwi.co.uk/arable/osr-growers-face-disaster-from-flea-](https://www.fwi.co.uk/arable/osr-growers-face-disaster-from-flea-beetle-attack)
383 [beetle-attack](https://www.fwi.co.uk/arable/osr-growers-face-disaster-from-flea-beetle-attack) [Accessed 10 Feb. 2020].

384 Clarke, J. et al. (2009) Pesticide availability for cereals and oilseeds following revision of
385 Directive 91/414/EEC; effects of losses and new research priorities.HGCA Research Review
386 70,ADAS, Boxworth, Cambridge CB23 4NN

387 Clarke, A. (2014). Insecticide resistance confirmed in UK flea beetles. *Farmers Weekly*.
388 [online] Available at: [https://www.fwi.co.uk/arable/insecticide-resistance-confirmed-in-uk-](https://www.fwi.co.uk/arable/insecticide-resistance-confirmed-in-uk-flea-beetles)
389 [flea-beetles](https://www.fwi.co.uk/arable/insecticide-resistance-confirmed-in-uk-flea-beetles) [Accessed 10 Feb. 2020].

390 Dewar, A. M. The adverse impact of the neonicotinoid seed treatment ban on crop protection
391 in oilseed rape in the United Kingdom Pest Manage. Sci. **2017**, 73 (7) 1305– 1309DOI:
392 10.1002/ps.4511

393 Dyer, A. (2019). Cabbage stem flea beetle losses threaten oilseed rape's place in the
394 rotation. *Farmers Guardian*. [online] Available at:
395 [https://www.fginsight.com/news/news/cabbage-stem-flea-beetle-losses-threaten-oilseed-](https://www.fginsight.com/news/news/cabbage-stem-flea-beetle-losses-threaten-oilseed-rape-place-in-the-rotation-84863)
396 [rape-place-in-the-rotation-84863](https://www.fginsight.com/news/news/cabbage-stem-flea-beetle-losses-threaten-oilseed-rape-place-in-the-rotation-84863) [Accessed 29 Jan. 2020].

397 EU Commission (2013) EU, L 139/12 Commission Implementing Regulation (EU) No.
398 485/2013, Official Journal of the European Union. 56: pp. 12–26.

399 EU Commission (2018) Neonicotinoids. Available at:
400 https://ec.europa.eu/food/plant/pesticides/approval_active_substances/approval_renewal/neon
401 [icotinoids_en](https://ec.europa.eu/food/plant/pesticides/approval_active_substances/approval_renewal/neon).

402 FarmingUK Team (2015). Neonicotinoid ban continuing to devastate oilseed rape crops.
403 *Farminguk*. [online] Available at: [https://www.farminguk.com/news/neonicotinoid-ban-](https://www.farminguk.com/news/neonicotinoid-ban-continuing-to-devastate-oilseed-rape-crops_35693.html)
404 [continuing-to-devastate-oilseed-rape-crops_35693.html](https://www.farminguk.com/news/neonicotinoid-ban-continuing-to-devastate-oilseed-rape-crops_35693.html) [Accessed 10 Feb. 2020].

405 FarmingUK Team (2017). Neonicotinoid ban cost farmers £18.4m in 2016, figures show.
406 *FarmingUK*. [online] Available at: [https://www.farminguk.com/news/neonicotinoid-ban-](https://www.farminguk.com/news/neonicotinoid-ban-cost-farmers-18-4m-in-2016-figures-show_46260.html)
407 [cost-farmers-18-4m-in-2016-figures-show_46260.html](https://www.farminguk.com/news/neonicotinoid-ban-cost-farmers-18-4m-in-2016-figures-show_46260.html) [Accessed 10 Feb. 2020].

408 Gillbard, E. (2019). Survey reveals true cost of flea beetle on OSR yields. *Farmers Weekly*.
409 [online] Available at: [https://www.fwi.co.uk/arable/harvest/survey-reveals-true-cost-of-flea-](https://www.fwi.co.uk/arable/harvest/survey-reveals-true-cost-of-flea-beetle-on-osr-yields)
410 [beetle-on-osr-yields](https://www.fwi.co.uk/arable/harvest/survey-reveals-true-cost-of-flea-beetle-on-osr-yields) [Accessed 29 Jan. 2020].

411 Graham, C. W. and Alford, D. V. (1981) 'The distribution and importance of cabbage stem
412 flea beetle (*Psylliodes chrysocephala* (L.)) on winter oilseed rape in England', *Plant*
413 *Pathology*, 30(3), pp. 141–145.

414 Heimbach, U. and Müller, A. (2013) 'Incidence of pyrethroid-resistant oilseed rape pests in
415 Germany', *Pest Management Science*. Wiley-Blackwell, 69(2), pp. 209–216.

416 Hill, O. (2017). 8% of British rapeseed crop lost to flea beetle and drought. *Farmers Weekly*.
417 [online] Available at: [https://www.fwi.co.uk/arable/osr/8-percent-of-british-rapeseed-crop-](https://www.fwi.co.uk/arable/osr/8-percent-of-british-rapeseed-crop-lost-to-flea-beetle-and-drought)
418 [lost-to-flea-beetle-and-drought](https://www.fwi.co.uk/arable/osr/8-percent-of-british-rapeseed-crop-lost-to-flea-beetle-and-drought) [Accessed 10 Feb. 2020].

419 Højland, D. H. and Kristensen, M. (2018) Target-site and metabolic resistance against λ -
420 cyhalothrin in cabbage stem flea beetles in Denmark, *Bulletin of Insectology*, 71(1), pp. 45–
421 49.

422 Højland, D. H., Nauen, R., Foster, S. P., Williamson, M. S. and Kristensen, M. (2015)
 423 Incidence, Spread and Mechanisms of Pyrethroid Resistance in European Populations of the
 424 Cabbage Stem Flea Beetle, *Psylliodes chrysocephala* L. (Coleoptera: Chrysomelidae). PLoS
 425 ONE 10(12): e0146045. doi:10.1371/journal.pone.0146045.

426 Livak, K.J., 1999. Allelic discrimination using fluorogenic probes and the 5' nuclease
 427 assay. *Genetic analysis: biomolecular engineering*, 14(5-6), pp.143-149.

428 Miller, A.L., Tindall, K. and Leonard, B.R. (2010) Bioassays for monitoring insecticide
 429 resistance. *Journal of visualized experiments: JoVE*, (46).

430 Nauen, R., Zimmer, C. T., Andrews, M., Slater, R., Bass, C., Ekbom, B., Gustafsson, G.,
 431 Hansen, L. M., Kristensen, M., Zebitz, C. P. W. and Williamson, M. S. (2012) 'Target-site
 432 resistance to pyrethroids in European populations of pollen beetle, *Meligethes aeneus* F.',
 433 *Pesticide Biochemistry and Physiology*, 103, pp. 173–180.

434 Slater, R., Ellis, S., Genay, J.P., Heimbach, U., Huart, G., Sarazin, M., Longhurst, C., Müller,
 435 A., Nauen, R., Rison, J.L. and Robin, F. (2011) Pyrethroid resistance monitoring in European
 436 populations of pollen beetle (*Meligethes* spp.): a coordinated approach through the Insecticide
 437 Resistance Action Committee (IRAC). *Pest management science*, 67(6), pp.633-638.

438 Williams, I. H. (2004) Advances in Insect Pest Management of Oilseed Rape in Europe, in
 439 *Insect Pest Management*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 181–208.

440 Wynn S, Ellis S and Alves L, *Cabbage Stem Flea Beetle Snapshot Assessment – September*
 441 *2014*. [Online]. Available: <http://cereals.ahdb.org.uk/media/507048/pr546-extension.pdf>.

442 Zimmer, C. T., Bass, C., Williamson, M. S., Kaussmann, M., Wölfel, K., Gutbrod, O. and
 443 Nauen, R. (2014) 'Molecular and functional characterization of CYP6BQ23, a cytochrome
 444 P450 conferring resistance to pyrethroids in European populations of pollen beetle,

445 *Meligethes aeneus*', *Insect Biochemistry and Molecular Biology*, 45(1), pp. 18–29.

446 Zimmer, C. T. and Nauen, R. (2011) Cytochrome P450 mediated pyrethroid resistance in
447 European populations of *Meligethes aeneus* (Coleoptera: Nitidulidae) *Pest. Biochem.*
448 *Physiol.*, 100 (2011), pp. 264-272.

449 Zimmer, C. T., Müller, A., Heimbach, U. and Nauen, R. (2014) 'Target-site resistance to
450 pyrethroid insecticides in German populations of the cabbage stem flea beetle, *Psylliodes*
451 *chrysocephala* L. (Coleoptera: Chrysomelidae)', *Pesticide Biochemistry and Physiology*,
452 108(1), pp. 1–7.